

Theoretical Mathematics & Applications, vol.2, no.3, 2012, 23-38
ISSN: 1792-9687 (print), 1792-9709 (online)
Scienpress Ltd, 2012

Hybrid descent-like halpern iteration methods for two nonexpansive mappings and semigroups on two sets

Nguyen Buong ¹and Nguyen Duc Lang²

Abstract

In this paper, we introduce some new iteration methods based on the hybrid method in mathematical programming, the descent-like iterative method and the Halpern's method for finding a common fixed point of two nonexpansive mappings and nonexpansive semigroups on two closed and convex subsets in Hilbert spaces.

Mathematics Subject Classification: 47J05, 47H09, 49J30

Keywords: Metric projection, Fixed point, Nonexpansive Mappings and Semigroups

¹ Vietnamese Academy of Science and Technology Institute of Information Technology
Technology 18, Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam.

² Viet Nam, Thainguyen University, University of Sciences,
e-mail: nguyenduclang2002@yahoo.com

1 Introduction

Let H be a real Hilbert space with the scalar product and the norm denoted by the symbols $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively, and let C be a nonempty, closed and convex subset of H . Denote by $P_C x$ the metric projection from $x \in H$ onto C . Let T be a nonexpansive mapping on C , i.e., $T : C \rightarrow C$ and $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. We use $F(T)$ to denote the set of fixed points of T , i.e., $F(T) = \{x \in C : x = Tx\}$. We know that $F(T)$ is nonempty, if C is bounded, for more details see [1].

Let $\{T(t) : t > 0\}$ be a nonexpansive semigroup on C , that is,

- (1) for each $t > 0$, $T(t)$ is a nonexpansive mapping on C ;
- (2) $T(0)x = x$ for all $x \in C$;
- (3) $T(t_1 + t_2) = T(t_1) \circ T(t_2)$ for all $t_1, t_2 > 0$; and
- (4) for each $x \in C$, the mapping $T(\cdot)x$ from $(0, \infty)$ into C is continuous.

Denote by $\mathcal{F} = \bigcap_{t>0} F(T(t))$ the set of common fixed points for the semigroup $\{T(t) : t > 0\}$. We know that \mathcal{F} is a closed convex subset in H and $\mathcal{F} \neq \emptyset$ if C is compact (see, [2]).

Let $C_i, i = 1, 2$, be two closed and convex subsets in H . Let T_i and $\{T_i(t) : t > 0\}$, $i = 1, 2$, be two nonexpansive mappings and semigroups on C_i , respectively. The problems studied in this paper is to find two elements

$$p \in F := F(T_1) \cap F(T_2) \tag{1.1}$$

and

$$q \in \mathcal{F}_{1,2} := \mathcal{F}_1 \cap \mathcal{F}_2, \tag{1.2}$$

where $\mathcal{F}_i = \bigcap_{t>0} F(T_i(t))$. Assume that F and $\mathcal{F}_{1,2}$ are not empty. Some particular cases of (1.1) and (1.2) are the following:

- (i) when $T_1 = T_2 = I$, the identity mapping in H , (1.1) is the convex feasibility problem studied in [3].
- (ii) when $C_1 = C_2 = C$, problems (1.1) and (1.2) are considered in [4]-[6].

For finding a fixed point of a nonexpansive mapping T on C , in 1953, Mann [7] proposed the following method:

$$\begin{aligned} x_0 &\in C \quad \text{any element,} \\ x_{n+1} &= \alpha_n x_n + (1 - \alpha_n) T x_n, \quad n \geq 0, \end{aligned} \tag{1.3}$$

that converges only weakly, in general (see [8] for an example). In 1967, Halpern [9] firstly proposed the following iteration process:

$$x_{n+1} = \beta_n u + (1 - \beta_n)Tx_n, \quad n \geq 0, \quad (1.4)$$

where u, x_0 are two fixed elements in C and $\{\beta_n\} \subset (0, 1)$. He pointed out that the conditions $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=0}^{\infty} \beta_n = \infty$ are necessary in the sense that, if the iteration (1.4) converges to a fixed point of T , then these conditions must be satisfied. Further, the iteration method was investigated by Lions [10], Reich [11], Wittmann [12] and Song [13]. Recently, Alber [14] proposed the following descent-like method

$$x_{n+1} = P_C(x_n - \mu_n[x_n - Tx_n]), n \geq 0, \quad (1.5)$$

and proved that if $\{\mu_n\} : \mu_n > 0, \mu_n \rightarrow 0$, as $n \rightarrow \infty$ and $\{x_n\}$ is bounded, then:

- (i) there exists a weak accumulation point $\tilde{x} \in C$ of $\{x_n\}$;
- (ii) all weak accumulation points of $\{x_n\}$ belong to $F(T)$; and
- (iii) if $F(T)$ is a singleton, i.e., $F(T) = \{\tilde{x}\}$, then $\{x_n\}$ converges weakly to \tilde{x} .

To obtain strong convergence for (1.3), Nakajo and Takahashi [15] introduced the hybrid Mann's iteration method:

$$\begin{aligned} x_0 &\in C, \\ y_n &= \alpha_n x_n + (1 - \alpha_n)Tx_n, \\ C_n &= \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, \end{aligned} \quad (1.6)$$

where $\{\alpha_n\} \subset [0, a]$ for some $a \in [0, 1)$. They showed that $\{x_n\}$ defined by (1.6) converges strongly to $P_{F(T)}x_0$ as $n \rightarrow \infty$. Recently, Yanes and Xu [16] adapted the iteration process (1.4) as follows:

$$\begin{aligned} x_0 &\in C \quad \text{any element,} \\ y_n &= \beta_n x_0 + (1 - \beta_n)Tx_n, \\ C_n &= \{z \in C : \|y_n - z\|^2 \leq \|x_n - z\|^2 \\ &\quad + \beta_n(\|x_0\|^2 + 2\langle x_n - x_0, z \rangle)\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0. \end{aligned} \quad (1.7)$$

They proved that if T is a nonexpansive mapping on a closed convex subset C with $F(T) \neq \emptyset$ and the sequence $\{\beta_n\} \subset (0, 1)$ is chosen such that $\lim_{n \rightarrow \infty} \beta_n = 0$, then the sequence $\{x_n\}$ defined by (1.7) converges strongly to $P_{F(T)}x_0$ as $n \rightarrow \infty$.

For finding an element $p \in \mathcal{F}$, Nakajo and Takahashi [15] also introduced an iteration procedure as follows:

$$\begin{aligned} x_0 &\in C \quad \text{any element,} \\ y_n &= \alpha_n x_n + (1 - \alpha_n) \frac{1}{t_n} \int_0^{t_n} T(s)x_n ds, \\ C_n &= \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, n \geq 0, \end{aligned} \tag{1.8}$$

where $\alpha_n \in [0, a]$ for some $a \in [0, 1)$ and $\{t_n\}$ is a positive real number divergent sequence. Under the conditions on $\{\alpha_n\}$ and $\{t_n\}$, the sequence $\{x_n\}$ defined by (1.8) converges strongly to $P_{\mathcal{F}}x_0$.

If $C \equiv H$, then C_n and Q_n in (1.6)-(1.8) are two halfspaces. So, the projection x_{n+1} onto $C_n \cap Q_n$ in these methods can be found by an explicit formula [17]. Clearly, if C is a proper subset of H , then C_n and Q_n in (1.6)-(1.8) are not two halfspaces. Then, the following problem is posed: how to construct the closed convex subsets C_n and Q_n and if we can express x_{n+1} of (1.6)-(1.8) in a similar form as in [17]? This problem is solved very recently in [18]-[20]. In this works, C_n and Q_n are replaced by two halfspaces and y_n is the right hand side of (1.5) with a modification. In this paper, motivated by (1.5), (1.7) and [14], [15], to solve problems (1.1) and (1.2) we introduce the following new iteration processes:

$$\begin{aligned} x_0 &\in H \quad \text{any element,} \\ z_n &= x_n - \mu_n(x_n - T_1 P_{C_1} x_n), \\ y_n &= \beta_n x_0 + (1 - \beta_n) T_2 P_{C_2} z_n, \\ H_n &= \{z \in H : \|y_n - z\|^2 \leq \|x_n - z\|^2 \\ &\quad + \beta_n(\|x_0\|^2 + 2\langle x_n - x_0, z \rangle)\}, \\ W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n} x_0, n \geq 0; \end{aligned} \tag{1.9}$$

and

$$\begin{aligned}
 x_0 &\in H \quad \text{any element,} \\
 z_n &= x_n - \mu_n \left(x_n - \frac{1}{t_n} \int_0^{t_n} T_1(s) P_{C_1} x_n ds \right), \\
 y_n &= \beta_n x_0 + (1 - \beta_n) \frac{1}{t_n} \int_0^{t_n} T_2(s) P_{C_2} z_n ds, \\
 H_n &= \{z \in H : \|y_n - z\|^2 \leq \|x_n - z\|^2 \\
 &\quad + \beta_n (\|x_0\|^2 + 2\langle x_n - x_0, z \rangle)\}, \\
 W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\
 x_{n+1} &= P_{H_n \cap W_n} x_0, \quad n \geq 0.
 \end{aligned} \tag{1.10}$$

We shall prove the strong convergence of the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ defined by (1.9) and (1.10) to some elements p and q in sections 2 and 3, respectively.

Below, the symbols \rightharpoonup and \rightarrow denote weak and strong convergence, respectively.

2 Strong convergence to a common fixed point of two nonexpansive mappings

We formulate the following facts needed in the proof of our results.

Lemma 2.1 [21]. *Let H be a real Hilbert. There holds the following identity: $\|\lambda x + (1 - \lambda)y\|^2 = \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 - \lambda(1 - \lambda)\|x - y\|^2$.*

Lemma 2.2 [16]. *Let C be a nonempty, closed and convex subset of a real Hilbert space H . For any $x \in H$, there exists a unique $z \in C$ such that $\|z - x\| \leq \|y - x\|$ for all $y \in C$, and $z = P_C x$ if and only if $\langle z - x, y - z \rangle \geq 0$ for all $y \in C$.*

Lemma 2.3. (Demiclosedness principle) [21]. *If C is a nonempty, closed and convex subset of a real Hilbert space H , T is a nonexpansive mapping on C , $\{x_n\}$ is a sequence in C such that $x_n \rightharpoonup x$ and $x_n - Tx_n \rightarrow 0$, then $x - Tx = 0$.*

Lemma 2.4 [22]. *Every Hilbert space H has Randon-Riesz property or Kadec-*

Klee property, that is, for a sequence $\{x_n\} \subset H$ with $x_n \rightharpoonup x$ and $\|x_n\| \rightarrow \|x\|$, then there holds $x_n \rightarrow x$.

Now, we are in a position to prove the following result.

Theorem 2.5. *Let C_1 and C_2 be two nonempty, closed and convex subsets in a real Hilbert space H and let T_1 and T_2 be two nonexpansive mappings on C_1 and C_2 , respectively, such that $F := F(T_1) \cap F(T_2) \neq \emptyset$. Assume that $\{\mu_n\}$ and $\{\beta_n\}$ are sequences in $[0,1]$ such that $\mu_n \in (a, b)$ for some $a, b \in (0, 1)$ and $\beta_n \rightarrow 0$. Then, the sequences $\{x_n\}, \{z_n\}$ and $\{y_n\}$, defined by (1.9), converge strongly to the same point $u_0 = P_F x_0$, as $n \rightarrow \infty$.*

Proof. First, note that

$$\|y_n - z\|^2 \leq \|x_n - z\|^2 + \beta_n(\|x_0\|^2 + 2\langle x_n - x_0, z \rangle)$$

is equivalent to

$$\langle (1 - \beta_n)x_n + \beta_n x_0 - y_n, z \rangle \leq \langle x_n - y_n, x_n \rangle - \frac{1}{2}\|y_n - x_n\|^2 + \frac{\beta_n}{2}\|x_0\|^2.$$

Thus, H_n is a halfspace. It is clear that $F(T) = F(TP_C) := \{p \in H : TP_C p = p\}$ for any mapping T from C into C . So, we have that $F = F(\tilde{T}_1) \cap F(\tilde{T}_2)$ where $\tilde{T}_i = T_i P_{C_i}$, $i = 1, 2$, and \tilde{T}_i , $i = 1, 2$, are also two nonexpansive mappings on H . Hence, by (1.9) and Lemma 2.1, we obtain for any $p \in F$ that

$$\begin{aligned} \|z_n - p\|^2 &= \|(1 - \mu_n)(x_n - p) + \mu_n(\tilde{T}_1 x_n - p)\|^2 \\ &= (1 - \mu_n)\|x_n - p\|^2 + \mu_n\|\tilde{T}_1 x_n - p\|^2 \\ &\quad - (1 - \mu_n)\mu_n\|x_n - \tilde{T}_1 x_n\|^2 \\ &\leq (1 - \mu_n)\|x_n - p\|^2 + \mu_n\|x_n - p\|^2 \\ &\quad - (1 - \mu_n)\mu_n\|x_n - \tilde{T}_1 x_n\|^2 \\ &\leq \|x_n - p\|^2 - (1 - \mu_n)\mu_n\|x_n - \tilde{T}_1 x_n\|^2 \leq \|x_n - p\|^2. \end{aligned} \tag{2.1}$$

By the similar argument and the convexity of $\|\cdot\|^2$, we also obtain

$$\begin{aligned}
 \|y_n - p\|^2 &= \|\beta_n x_0 + (1 - \beta_n) \tilde{T}_2 z_n - p\|^2 \\
 &\leq \beta_n \|x_0 - p\|^2 + (1 - \beta_n) \|\tilde{T}_2 z_n - \tilde{T}_2 p\|^2 \\
 &\leq \beta_n \|x_0 - p\|^2 + (1 - \beta_n) \|z_n - p\|^2 \\
 &\leq \beta_n \|x_0 - p\|^2 + (1 - \beta_n) \|x_n - p\|^2 \\
 &= \|x_n - p\|^2 + \beta_n (\|x_0 - p\|^2 - \|x_n - p\|^2) \\
 &= \|x_n - p\|^2 + \beta_n (\|x_0\|^2 + 2\langle x_n - x_0, p \rangle).
 \end{aligned}$$

Therefore, $p \in H_n$ for all $n \geq 0$. It means that $F(T) \subset H_n$ for all $n \geq 0$.

Next, we show by mathematical induction that $F(T) \subset H_n \cap W_n$ for each $n \geq 0$. For $n = 0$, we have $W_0 = H$, and hence $F(T) \subset H_0 \cap W_0$. Suppose that x_i is given and $F(T) \subset H_i \cap W_i$ for some $i > 0$. There exists a unique element $x_{i+1} \in H_i \cap W_i$ such that $x_{i+1} = P_{H_i \cap W_i} x_0$. Therefore, by Lemma 2.2,

$$\langle x_{i+1} - x_0, p - x_{i+1} \rangle \geq 0$$

for each $p \in H_i \cap W_i$. Since $F(T) \subset H_i \cap W_i$, we get $F(T) \subset W_{i+1}$. So, we have $F(T) \subset H_{i+1} \cap W_{i+1}$.

Further, since $F(T)$ is a nonempty, closed and convex subset of H , there exists a unique element $u_0 \in F(T)$ such that $u_0 = P_{F(T)} x_0$. From $x_{n+1} = P_{H_n \cap W_n}(x_0)$, we obtain

$$\|x_{n+1} - x_0\| \leq \|z - x_0\|$$

for every $z \in H_n \cap W_n$. As $u_0 \in F(T) \subset W_n$, we get

$$\|x_{n+1} - x_0\| \leq \|u_0 - x_0\| \quad \forall n \geq 0. \quad (2.2)$$

This implies that $\{x_n\}$ is bounded. Now, we show that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (2.3)$$

From the definition of W_n and Lemma 2.2, we have $x_n = P_{W_n} x_0$. As $x_{n+1} \in H_n \cap W_n$, we obtain

$$\|x_{n+1} - x_0\| \geq \|x_n - x_0\| \quad \forall n \geq 0.$$

Therefore, $\{\|x_n - x_0\|\}$ is a nondecreasing and bounded sequence. So, there exists $\lim_{n \rightarrow \infty} \|x_n - x_0\| = c$. On the other hand, from $x_{n+1} \in W_n$, it follows that

$$\langle x_n - x_0, x_{n+1} - x_n \rangle \geq 0,$$

and hence

$$\begin{aligned}\|x_n - x_{n+1}\|^2 &= \|x_n - x_0 - (x_{n+1} - x_0)\|^2 \\ &= \|x_n - x_0\|^2 - 2\langle x_n - x_0, x_{n+1} - x_0 \rangle + \|x_{n+1} - x_0\|^2 \\ &\leq \|x_{n+1} - x_0\|^2 - \|x_n - x_0\|^2 \quad \forall n \geq 0.\end{aligned}$$

Thus, (2.3) is proved by using the last inequality and $\lim_{n \rightarrow \infty} \|x_n - x_0\| = c$.

Next, since $x_{n+1} \in H_n$ we have that

$$\|y_n - x_{n+1}\|^2 \leq \|x_n - x_{n+1}\|^2 + \beta_n(\|x_0\| + 2\langle x_n - x_0, z \rangle).$$

Therefore, from (2.3), the boundness of $\{x_n\}$, $\beta_n \rightarrow 0$ and the last inequality, it follows that

$$\lim_{n \rightarrow \infty} \|y_n - x_{n+1}\| = 0. \quad (2.4)$$

This together with (2.3) implies that

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \quad (2.5)$$

Noticing that $\tilde{T}_2 z_n = y_n - \beta_n(x_n - \tilde{T}_2 z_n) + \beta_n(x_n - x_0)$, we have

$$\|x_n - \tilde{T}_2 z_n\| \leq \|x_n - y_n\| + \beta_n \|x_n - \tilde{T}_2 z_n\| + \beta_n \|x_n - x_0\|.$$

From (2.2) and the last inequality, it follows that

$$\|x_n - \tilde{T}_2 z_n\| \leq \frac{1}{1 - \beta_n} \left(\|x_n - y_n\| + \beta_n \|x_0 - x_0\| \right).$$

By $\beta_n \rightarrow 0$ ($\beta_n \leq 1 - \beta$ for some $\beta \in (0, 1)$), (2.5) and the last inequality, we obtain

$$\lim_{n \rightarrow \infty} \|x_n - \tilde{T}_2 z_n\| = 0. \quad (2.6)$$

Now, we shall prove that $\|x_n - \tilde{T}_1 x_n\| \rightarrow 0$ and $\|x_n - \tilde{T}_2 x_n\| \rightarrow 0$, as $n \rightarrow \infty$. Indeed, since $\{x_n\}$ is bounded, for any $p \in F$ and any subsequence $\{\tilde{T}_1 x_{n_k} - x_{n_k}\}$ of $\{\tilde{T}_1 x_n - x_n\}$ there exists a subsequence $\{x_{n_j}\} \subset \{x_{n_k}\}$ such that

$$\lim_{j \rightarrow \infty} \|x_{n_j} - p\| = \limsup_{k \rightarrow \infty} \|x_{n_k} - p\| = a.$$

By (2.6), (2.1) and the following inequalities

$$\begin{aligned}\|x_{n_j} - p\| &\leq \|x_{n_j} - \tilde{T}_2 z_{n_j}\| + \|\tilde{T}_2 z_{n_j} - p\| \\ &\leq \|x_{n_j} - \tilde{T}_2 z_{n_j}\| + \|z_{n_j} - p\| \\ &\leq \|x_{n_j} - \tilde{T}_2 z_{n_j}\| + \|x_{n_j} - p\|,\end{aligned}$$

we get that

$$\lim_{j \rightarrow \infty} \|x_{n_j} - p\| = \lim_{j \rightarrow \infty} \|z_{n_j} - p\| = a.$$

Again from (2.1) and the condition on μ_n , it implies that

$$a(1-b)\|\tilde{T}_1 x_{n_j} - x_{n_j}\| \leq \|x_{n_j} - p\| - \|z_{n_j} - p\|.$$

So, $\|\tilde{T}_1 x_{n_j} - x_{n_j}\| \rightarrow 0$ and hence $\|\tilde{T}_1 x_n - x_n\| \rightarrow 0$, as $n \rightarrow \infty$. Further, since

$$\begin{aligned} \|\tilde{T}_2 x_n - x_n\| &\leq \|\tilde{T}_2 x_n - \tilde{T}_2 z_n\| + \|\tilde{T}_2 z_n - x_n\| \\ &\leq \|x_n - z_n\| + \|\tilde{T}_2 z_n - x_n\|, \end{aligned}$$

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = \lim_{n \rightarrow \infty} \mu_n \|\tilde{T}_1 x_n - x_n\| = 0, \quad (2.7)$$

by (2.6) and $\|\tilde{T}_1 x_n - x_n\| \rightarrow 0$, we also obtain that $\|\tilde{T}_2 x_n - x_n\| \rightarrow 0$. Since $\{x_n\}$ is bounded, there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ that converges weakly to some element $p \in H$ as $i \rightarrow \infty$. By Lemmas 2.3 and $\|\tilde{T}_1 x_n - x_n\|, \|\tilde{T}_2 x_n - x_n\| \rightarrow 0$, we have that $p \in F$.

Now, from (2.2) and the weak lower semicontinuity of the norm it implies that

$$\|x_0 - u_0\| \leq \|x_0 - p\| \leq \liminf_{j \rightarrow \infty} \|x_0 - x_{n_j}\| \leq \limsup_{j \rightarrow \infty} \|x_0 - x_{n_j}\| \leq \|x_0 - u_0\|.$$

Thus, we obtain $\lim_{j \rightarrow \infty} \|x_0 - x_{n_j}\| = \|x_0 - u_0\| = \|x_0 - p\|$. This implies $x_{k_j} \rightarrow p = u_0$ by Lemma 2.4. By the uniqueness of the projection $u_0 = P_F x_0$, we have that $x_n \rightarrow u_0$. Consequently, from (2.7) it follows that $z_n \rightarrow u_0$. From (2.5), we also get that $y_n \rightarrow u_0$. This completes the proof. \square

We have the following corollaries.

Corollary 2.6. *Let $C_i, i = 1, 2$, be two nonempty, closed and convex subsets in a real Hilbert space H . Let $T_i, i = 1, 2$, be two nonexpansive mappings on C_i such that $F(T_1) \cap F(T_2) \neq \emptyset$. Assume that $\{\mu_n\}$ is a sequence such that $0 < a \leq \mu_n \leq b < 1$. Then, the sequences $\{x_n\}$ and $\{y_n\}$, defined by*

$$\begin{aligned} x_0 &\in H \quad \text{any element,} \\ y_n &= T_2 P_{C_2}(x_n - \mu_n(x_n - T_1 P_{C_1} x_n)), \\ H_n &= \{z \in H : \|y_n - z\| \leq \|x_n - z\|\}, \\ W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n}(x_0), n \geq 0, \end{aligned}$$

converge strongly to the same point $u_0 = P_{F(T)}x_0$, as $n \rightarrow \infty$.

Proof. By putting $\beta_n \equiv 0$ in Theorem 2.5, we obtain the conclusion. \square

Corollary 2.7. *Let $C_i, i = 1, 2$, be two nonempty, closed and convex subsets in a real Hilbert space H such that $C := C_1 \cap C_2 \neq \emptyset$. Assume that $\{\mu_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$ such that $\mu_n \in (a, b)$ for some $a, b \in (0, 1)$ and $\beta_n \rightarrow 0$. Then, the sequences $\{x_n\}, \{z_n\}$ and $\{y_n\}$, defined by*

$$\begin{aligned} x_0 &\in H \quad \text{any element,} \\ z_n &= x_n - \mu_n(x_n - P_{C_1}x_n), \\ y_n &= \beta_n x_0 + (1 - \beta_n)P_{C_2}z_n, \\ H_n &= \{z \in H : \|y_n - z\|^2 \leq \|x_n - z\|^2 \\ &\quad + \beta_n(\|x_0\| + 2\langle x_n - x_0, z \rangle)\}, \\ W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n}x_0, n \geq 0, \end{aligned}$$

converge strongly to the same point $u_0 = P_C x_0$, as $n \rightarrow \infty$.

Proof. By putting $T_1 = T_2 = I$ in Theorem 2.5, we obtain the conclusion. \square

3 Strong convergence to a common fixed point of two nonexpansive semigroups

We need the following Lemma in the proof of our result.

Lemma 3.1 [23]. *Let C be a nonempty bounded closed convex subset in a real Hilbert space H and let $\{T(t) : t > 0\}$ be a nonexpansive semigroup on C . Then, for any $h > 0$*

$$\limsup_{t \rightarrow \infty} \sup_{y \in C} \left\| T(h) \left(\frac{1}{t} \int_0^t T(s)y ds \right) - \frac{1}{t} \int_0^t T(s)y ds \right\| = 0.$$

Now, we prove the following result.

Theorem 3.2. *Let C_1 and C_2 be two nonempty closed convex subsets in a real Hilbert space H and let $\{T_1(t) : t > 0\}$ and $\{T_2(t) : t > 0\}$ be two nonexpansive semigroups on C_1 and C_2 , respectively, such that $\mathcal{F} = \mathcal{F}_1 \cap \mathcal{F}_2 \neq \emptyset$ where*

$\mathcal{F}_i = \cap_{t>0} F(T_i(t))$, $i = 1, 2$. Assume that $\{\mu_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$ such that $\mu_n \in (a, b)$ for some $a, b \in (0, 1)$ and $\beta_n \rightarrow 0$ and $\{t_n\}$ is a positive real divergent sequence. Then, the sequences $\{x_n\}$, $\{z_n\}$ and $\{y_n\}$, defined by (1.10), converge strongly to the same point $u_0 = P_{\mathcal{F}}x_0$, as $n \rightarrow \infty$.

Proof. For each $p \in \mathcal{F}$, we have for each $s > 0$ that

$$p = P_{C_i}p = \tilde{T}_i(s)p, \quad i = 1, 2,$$

where $\tilde{T}_i(s) = T_i(s)P_{C_i}$, and hence from (1.10) and Lemma 2.1, we obtain that

$$\begin{aligned} \|z_n - p\|^2 &= \left\| (1 - \mu_n)(x_n - p) + \mu_n \left(\frac{1}{t_n} \int_0^{t_n} \tilde{T}_1(s)x_n ds - p \right) \right\|^2 \\ &= \left\| (1 - \mu_n)(x_n - p) + \mu_n \left(\frac{1}{t_n} \int_0^{t_n} [\tilde{T}_1(s)x_n - \tilde{T}_1(s)p] ds \right) \right\|^2 \\ &= (1 - \mu_n)\|x_n - p\|^2 + \mu_n \left\| \frac{1}{t_n} \int_0^{t_n} \tilde{T}_1(s)x_n - \tilde{T}_1(s)p ds \right\|^2 \\ &\quad - (1 - \mu_n)\mu_n \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_1(s)x_n ds \right\|^2 \\ &\leq \|x_n - p\|^2 - (1 - \mu_n)\mu_n \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_1(s)x_n ds \right\|^2 \\ &\leq \|x_n - p\|^2. \end{aligned} \tag{3.1}$$

By the similar argument and the convexity of $\|\cdot\|^2$, we also obtain

$$\begin{aligned} \|y_n - p\|^2 &= \left\| \beta_n(x_0 - p) + (1 - \beta_n) \left(\frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s)z_n ds - p \right) \right\|^2 \\ &\leq \beta_n\|x_0 - p\|^2 + (1 - \beta_n) \left\| \frac{1}{t_n} \int_0^{t_n} [\tilde{T}_2(s)z_n - \tilde{T}_2(s)p] ds \right\|^2 \\ &\leq \beta_n\|x_0 - p\|^2 + (1 - \beta_n)\|z_n - p\|^2 \\ &\leq \beta_n\|x_0 - p\|^2 + (1 - \beta_n)\|x_n - p\|^2 \\ &= \|x_n - p\|^2 + \beta_n(\|x_0 - p\|^2 - \|x_n - p\|^2) \\ &= \|x_n - p\|^2 + \beta_n(\|x_0\|^2 + 2\langle x_n - x_0, p \rangle). \end{aligned}$$

Therefore, $p \in H_n$ for $n \geq 0$. It means that $\mathcal{F} \subset H_n$ for $n \geq 0$. As in the proof of Theorem 2.5, we can obtain the following properties:

(i) $\mathcal{F} \subset H_n \cap W_n$,

$$\|x_{n+1} - x_0\| \leq \|u_0 - x_0\|, u_0 = P_{\mathcal{F}}x_0 \tag{3.2}$$

for $n \geq 0$. This implies that $\{x_n\}$ is bounded.

(ii)

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (3.3)$$

$$\lim_{n \rightarrow \infty} \|y_n - x_{n+1}\| = 0. \quad (3.4)$$

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \quad (3.5)$$

Noticing that

$$\frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds = y_n - \beta_n \left(x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds \right) + \beta_n (x_n - x_0),$$

we have

$$\begin{aligned} \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds \right\| &\leq \|x_n - y_n\| \\ &+ \beta_n \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds \right\| + \beta_n \|x_n - x_0\|. \end{aligned}$$

From (3.2) and the last inequality, it follows that

$$\left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds \right\| \leq \frac{1}{1 - \beta_n} \left(\|x_n - y_n\| + \beta_n \|x_n - x_0\| \right).$$

By $\beta_n \rightarrow 0$ ($\beta_n \leq 1 - \beta$ for some $\beta \in (0, 1)$), (3.5) and the last inequality, we obtain

$$\lim_{n \rightarrow \infty} \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_2(s) z_n ds \right\| = 0. \quad (3.6)$$

As in the proof of Theorem 2.5, by using (3.6) we can obtain that

$$\lim_{n \rightarrow \infty} \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \right\| = 0, i = 1, 2, \quad (3.7)$$

and

$$\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0. \quad (3.8)$$

Since

$$\frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \in C_i, i = 1, 2,$$

we have that

$$\begin{aligned} \left\| P_{C_i} x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \right\| &= \left\| P_{C_i} x_n - P_{C_i} \frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \right\| \\ &\leq \left\| x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \right\|, \end{aligned}$$

and hence from (3.7) it implies that

$$\lim_{n \rightarrow \infty} \left\| P_{C_i} x_n - \frac{1}{t_n} \int_0^{t_n} \tilde{T}_i(s) x_n ds \right\| = 0, i = 1, 2. \quad (3.9)$$

Since $\{x_n\}$ is bounded, there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ that converges weakly to some element $q \in H$ as $j \rightarrow \infty$. From (3.7) and (3.9), we also obtain that $u_{n_j}^i := P_{C_i} x_{n_j} \rightarrow q$ as $j \rightarrow \infty$. It means that $q \in C_1 \cap C_2$. Then, for each $h > 0$, we have that

$$\begin{aligned} \|T_i(h)u_n^i - u_n^i\| &\leq \left\| T_i(h)u_n^i - T_i(h) \left(\frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds \right) \right\| \\ &\quad + \left\| T_i(h) \left(\frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds \right) - \frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds \right\| \\ &\quad + \left\| \frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds - u_n^i \right\| \\ &\leq 2 \left\| \frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds - u_n^i \right\| \\ &\quad + \left\| T(h) \left(\frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds \right) - \frac{1}{t_n} \int_0^{t_n} T_i(s)u_n^i ds \right\|. \end{aligned} \quad (3.10)$$

Let $C_0^i = \{z \in C_i : \|z - u_0\| \leq 2\|x_0 - u_0\|\}$. Since $u_0 = P_{\mathcal{F}}x_0 \in C_i$, we have that

$$\|u_{n_j}^i - u_0\| = \|P_{C_i}x_{n_j} - P_{C_i}u_0\| \leq \|x_{n_j} - u_0\| \leq 2\|x_0 - u_0\|.$$

So, C_0^i is a nonempty bounded closed convex subset. It is easy to verify that $\{T_i(t) : t > 0\}$ is a nonexpansive semigroup on C_0^i . By Lemma 3.1, we get

$$\lim_{n \rightarrow \infty} \left\| T_i(h) \left(\frac{1}{t_n} \int_0^{t_n} T(s)u_n^i ds \right) - \frac{1}{t_n} \int_0^{t_n} T(s)u_n^i ds \right\| = 0$$

for every fixed $h > 0$ and hence by (3.9)-(3.10) we obtain that

$$\lim_{j \rightarrow \infty} \|T_i(h)u_{n_j}^i - u_{n_j}^i\| = 0$$

for each $h > 0$. By Lemma 2.3, $q \in F(T_i(h))$ for all $h > 0$. It means that $q \in \mathcal{F}$. As in the proof of Theorem 2.5, by using (3.2), (3.5) and (3.8), we also obtain that the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$, defined by (1.10), converge strongly to u_0 as $n \rightarrow \infty$. This completes the proof. \square

Corollary 3.3. *Let C be a nonempty closed convex subset in a real Hilbert*

space H and let $\{T(t) : t > 0\}$ be a nonexpansive semigroup on C such that $\mathcal{F} = \bigcap_{t>0} F(T(t)) \neq \emptyset$. Assume that $\{\beta_n\}$ is a sequence in $[0,1]$ such that $\beta_n \rightarrow 0$. Then, the sequences $\{x_n\}$ and $\{y_n\}$, defined by

$$\begin{aligned} x_0 &\in H \quad \text{any element,} \\ y_n &= \beta_n x_0 + (1 - \beta_n) \frac{1}{t_n} \int_0^{t_n} T(s) P_C(x_n) ds, \\ H_n &= \{z \in H : \|y_n - z\|^2 \leq \|x_n - z\|^2 \\ &\quad + \beta_n (\|x_0\| + 2\langle x_n - x_0, z \rangle)\}, \\ W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n}(x_0), n \geq 0, \end{aligned}$$

converge strongly to the same point $u_0 = P_{\mathcal{F}}x_0$, as $n \rightarrow \infty$.

Proof. By putting $T_1(s) = I$ for all $s > 0$, $C_1 = H$, $C_2 = C$ and $T_2(s) = T(s)$ in Theorem 3.2, we obtain the conclusion. \square

Corollary 3.4. *Let C be a nonempty closed convex subset in a real Hilbert space H and let $\{T(t) : t > 0\}$ be a nonexpansive semigroup on C such that $\mathcal{F} = \bigcap_{t>0} F(T(t)) \neq \emptyset$. Assume that $\{\alpha_n\}$ is a sequence in $[0,1]$ such that $\alpha_n \rightarrow 1$. Then, the sequences $\{x_n\}$ and $\{y_n\}$, defined by*

$$\begin{aligned} x_0 &\in H \quad \text{any element,} \\ y_n &= \frac{1}{t_n} \int_0^{t_n} T(s) P_C \left(x_n - \mu_n \left[x_n - \frac{1}{t_n} \int_0^{t_n} T(s) P_C x_n ds \right] ds \right), \\ H_n &= \{z \in H : \|y_n - z\| \leq \|x_n - z\|\}, \\ W_n &= \{z \in H : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n}(x_0), n \geq 0, \end{aligned}$$

converge strongly to the same point $u_0 = P_{\mathcal{F}}x_0$, as $n \rightarrow \infty$.

Proof. By putting $\beta_n \equiv 0$, $C_2 = H$, $C_1 = C$, $T_2(s) = I$ and $T_1(s) = T(s)$ for all $s > 0$ in Theorem 3.2, we obtain the conclusion. \square

References

- [1] E.F. Browder, Fixed-point theorems for noncompact mappings in Hilbert spaces, *Proceed. Nat. Acad. Sci. USA*, **53**, (1965), 1272-1276.

- [2] R. DeMarr, Common fixed points for commuting contraction mappings, *Pacific J. Math.*, **13**, (1963), 1139-1141.
- [3] H.H. Bauschke, P.L. Combettes and D.R. Luke, A strong convergent reflection method for finding the projection onto the intersection of two closed convex sets in a Hilbert spaces, *J. of Approx. Theory*, **141**, (2006), 63-69.
- [4] S.H. Khan and H. Fukhar-ud-din, Weak and strong convergence of a scheme with errors for two nonexpansive mappings, *Nonl. Anal.*, **61**, (2005), 1295-1301.
- [5] L. Wei, Y.J. Cho and H. Zhou, A strong convergence theorem for common fixed points of two relative nonexpansive mappings and its applications, *J. Appl. Math. Comput.*, DOI 10.1007/s12190-008-0092-x.
- [6] K. Wattanawitoon and P. Kummam, Convergence theorems of modified Ishikawa iterative scheme for two nonexpansive semigroups, *Fixed point theory and applications*, **2010**, Article ID 914702, 12 pages.
- [7] W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.*, **4**, (1953), 506-510.
- [8] A. Genel and J. Lindenstrass, An example concerning fixed points, *Israel J. Math.*, **22**, (1975), 81-86.
- [9] B. Halpern, Fixed points of nonexpanding maps, *Bull. Am.Math. Soc.*, **73**, (1967), 957-961.
- [10] P.L. Lions, Approximation de points fixes de contractions, *C.R. Acad. Sci. Sér A-B Paris*, **284**, (1977), 1357-1359.
- [11] S. Reich, Strong convergence theorem for resolvents of accretive operators in Banach spaces, *J. Math. Anal. Appl.*, **75**, (1980), 287-292.
- [12] R. Wittmann, Approximation of fixed points of nonexpansive mappings, *Arch. Math.*, **59**, (1992), 486-491.
- [13] Y. Song, A new sufficient condition for strong convergence of Halpern type iterations, *Appl. Math. Comput.*, **198**(1-2), (2007), 721-728.

- [14] Ya.I. Alber, On the stability of iterative approximations to fixed points of nonexpansive mappings, *J. Math. Anal. Appl.*, **328**, (2007), 958-971.
- [15] K. Nakajo and W. Takahashi, Strong convergence theorem for nonexpansive mappings and nonexpansive semigroup, *J. Math. Anal. Applic.*, **279**, (2003), 372-379.
- [16] C. Martinez-Yanes and H.K. Xu, Strong convergence of the CQ method for fixed iteration processes, *Nonl. Anal.*, **64**, (2006), 2400-2411.
- [17] M.V. Solodov and B.F. Svaiter, Forcing strong convergence of proximal point iterations in Hilbert space, *Math. Progr.*, **87**, (2000), 189-202.
- [18] Nguyen Buong, Strong convergence theorem for nonexpansive semigroup in Hilbert space, *Nonl. Anal.*, **72**(12), (2010), 4534-4540.
- [19] Nguyen Buong, Strong convergence theorem of an iterative method for variational inequalities and fixed point problems in Hilbert spaces, *Applied Math. and Comp.*, **217**, (2010), 322-329.
- [20] Nguyen Buong and Nguyen Duc Lang, Hybrid Mann-Halpern iteration methods for nonexpansive mappings and semigroups, *Applied Math. and Comp.*, **218**, (2011), 2459-2466.
- [21] G. Marino and H.K. Xu, Weak and strong convergence theorems for strict pseudo-contractions in Hilbert spaces, *J. Math. Anal. Appl.*, **329**, (2007), 336-346.
- [22] K. Goebel and W.A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge Studies in Advanced Math., **28**, Cambridge Univ. Press, Cambridge, 1990.
- [23] T. Shimizu and W. Takahashi, Strong convergence to common fixed points of families of nonexpansive mappings, *J. Math. Anal. Appl.*, **211**, (1997), 71-83.